

Regulated oxygen sensing by protein hydroxylation in renal erythropoietin-producing cells

Roland H. Wenger and David Hoogewijs

Institute of Physiology and Zürich Center for Integrative Human Physiology ZIHP, University of Zürich, Zürich, Switzerland

Submitted 23 December 2009; accepted in final form 9 March 2010

Wenger RH, Hoogewijs D. Regulated oxygen sensing by protein hydroxylation in renal erythropoietin-producing cells. *Am J Physiol Renal Physiol* 298: F1287–F1296, 2010. First published March 10, 2010; doi:10.1152/ajprenal.00736.2009.—The kidney is a major site of systemic oxygen sensing, regulating blood erythrocyte and hence oxygen content by hypoxia-inducible erythropoietin (Epo) expression. A constant ratio between blood perfusion and oxygen consumption, a stable corticomedullary oxygen gradient, and a relatively low tissue PO_2 are the prerequisites for the function of renal Epo-producing and oxygen-sensing (REPOS) cells, which are located in the juxtamedullary cortex. In kidney disease, renal oxygen consumption is decreased, leading to an increase in PO_2 , dysfunction of REPOS cells, and anemia. The molecular principles of cellular oxygen sensing have been elucidated in the last few years, and genetically altered mouse models as well as hereditary diseases causing erythrocytosis have clarified the oxygen-signaling cascade leading to increased Epo expression in REPOS cells. However, the consequences of a number of recently discovered factors for the regulation of oxygen signaling in REPOS cells are unclear, asking for novel cell culture models which might be hampered by the putative neuron-like nature of this enigmatic cell type.

hypoxia; anemia; erythrocytosis; gene expression; kidney disease; Krebs cycle; reactive oxygen species; von Hippel-Lindau

TWO MAJOR ORGANS IN THE ADULT body measure the systemic arterial oxygenation of the circulating blood: the carotid body and the kidney. The carotid body is located close to the lung circulation and obtains freshly oxygenated blood. Its type I chemoreceptor cells reside in a highly perfused tissue, and their oxygenation depends largely on the oxygen partial pressure (PO_2) rather than the oxygen content of the blood. The carotid body thus serves as a feedback control system that acutely surveys the function of the lung. Every drop in PO_2 will immediately result in increased alveolar ventilation to normalize arterial PO_2 .

The kidney, on the other hand, features a unique oxygen gradient which is stabilized by the countercurrent exchange of oxygen along the corticomedullary axis, keeping tissue PO_2 at relatively low values (116). This oxygen gradient is independent of the relatively high renal perfusion (20% of cardiac output) since every change in perfusion will lead to a similar change in renal oxygen consumption. Contrary to most other organs, the relatively low renal oxygen consumption (only 7% of the oxygen is extracted from the blood) is a function of renal perfusion and not vice versa. The more blood flow, the higher the glomerular filtration rate, the more sodium reabsorption and hence oxygen requirement, but also the more oxygen the kidney obtains, leading to a stable relationship between oxygen supply and oxygen consumption.



The renal erythropoietin (Epo)-producing and oxygen-sensing cells, which we suggest to name “REPOS” cells, are peritubular, interstitial, fibroblast-like cells located along the corticomedullary oxygen gradient in the juxtamedullary cortex (7, 68, 70, 84, 126). Tissue PO_2 in this region depends on oxygen diffusion and consumption on its way from arterial blood to the REPOS cells. Because a constant fraction of oxygen is consumed already in the outer cortex, the dropping PO_2 is dependent on the oxygen loaded on hemoglobin according to the sigmoidal oxygen binding properties of hemoglobin. Therefore, oxygenation of REPOS cells depends largely on the regional blood oxygen content rather than the arterial PO_2 . As illustrated in Fig. 1A, a drop in oxygen content, maybe due to inspiratory hypoxia or anemia, will lead to an increase in the number of REPOS cells that detectably produce Epo, extending their recruitment to the border of the subcapsular tissue (30, 69). These REPOS cells account for ~90% of total Epo synthesis in the adult (67). Following endocrine secretion, Epo stimulates bone marrow erythropoiesis to counteract decreased oxygen content by increased oxygen transport capacity (58).

Whereas the molecular oxygen-sensing mechanisms in the carotid body remain enigmatic, the last decade brought a wealth of novel insights, but also novel questions and novel therapeutic opportunities, on the oxygen-sensing and -signaling pathways regulating renal Epo synthesis.

Ubiquitous Cellular Oxygen Sensing by Protein Hydroxylation

The molecular mechanisms originally identified in hepatocytes and REPOS cells apply to all cells of the body: prolyl-

Address for reprint requests and other correspondence: R. H. Wenger, Institute of Physiology, Univ. of Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland (e-mail: roland.wenger@access.uzh.ch).



4-hydroxylase domain (PHD) enzymes continuously “sense” the oxygen levels and covalently modify hypoxia-inducible transcription factor- α (HIF- α) subunits (62, 123, 141). In the presence of sufficient oxygen, two distinct HIF- α prolyl residues within an oxygen-dependent degradation domain are hydroxylated in a nonreversible reaction involving oxygen, 2-oxoglutarate, ferrous iron, and probably vitamin C (54, 56). One oxygen atom is used for prolyl-4-hydroxylation, the other for oxidative decarboxylation of 2-oxoglutarate to form succinate and CO₂. Vitamin C might be involved in keeping the central iron atom in its ferrous form, but other reducing agents apparently can substitute for vitamin C (unpublished observations). This “intrinsic” reduction-oxidation (redox) reaction might be sensitive to externally generated reactive oxygen species (ROS), e.g., by mitochondrial electron transport chain complexes or by NADPH oxidases (2), but the physiological relevance of these “extrinsic” pathways is currently unknown. However, the entire PHD-HIF-von Hippel-Lindau (VHL) system can be reconstituted in a cell-free system in vitro where it shows the same oxygen-dependent HIF- α hydroxylation characteristics in the absence of any ROS-generating enzymes (146), suggesting that the intrinsic redox reaction is both necessary and sufficient for cellular oxygen sensing.

Under normoxic conditions, hydroxylation of HIF- α increases the affinity for binding by the tumor suppressor protein VHL, which serves as a recognition interface for an ubiquitin E3 ligase complex (85). Polyubiquitylation and destruction in proteasomes is a very rapid process that can be reverted instantaneously if oxygen supply is ceased (59). Under hypoxic conditions, HIF- α remains stable, heterodimerizes with HIF- β , and transcriptionally activates a large number of genes involved in the adaptation to decreased oxygen supply, including the gene encoding Epo (144). Another oxygen-dependent hydroxylase, factor-inhibiting HIF (FIH), triggers the transcriptional activity of HIF by asparagine hydroxylation, which blocks the recruitment of transcriptional coactivators (72, 76).

Hereditary Erythrocytosis Reveals PHD2, VHL, and HIF-2 α as Key Players in Renal Oxygen-Regulated Epo Gene Expression

In vertebrates, both the PHD oxygen sensors and the HIF- α transcription factor subunits are encoded by three distinct but related genes. The PHD family is encoded by the human *EGLN2*, *EGLN1*, and *EGLN3* genes and their respective gene products are called PHD1, PHD2, PHD3, or HIF prolyl hydroxylase (HPH) HPH3, HPH2, and HPH1 (20, 31). All three family members are expressed in the kidney (122). The physiological relevance of a fourth family member, called PH-4, is less clear: HIF- α seems to be regulated under PH-4 overexpression conditions only (66, 96). The human *HIF1A*, *HIF2A*, and *HIF3A* genes encode HIF-1 α , HIF-2 α , and HIF-3 α , respectively. There exist two related genes even for HIF- β , also known as aryl hydrocarbon receptor nuclear translocator (ARNT)-1 and ARNT-2, but only HIF- β /ARNT seems to serve

as an HIF- α heterodimerization partner. While HIF-1 and HIF-2 $\alpha\beta$ heterodimers function as transcriptional activators of oxygen-regulated target genes, the role of HIF-3 α is less clear, and a short splice variant of HIF-3 α , termed inhibitory PAS protein (IPAS), functions as a hypoxia-inducible transcriptional repressor (77). The HIF family has been discovered based on the cloning of HIF-1 α and HIF- β /ARNT by virtue of their binding to the 3' hypoxia response element (HRE) of the Epo gene in hepatoma cell lines (139). Thus it came as a surprise that it is not HIF-1 which regulates renal Epo expression. Mouse gene targeting and human hereditary diseases provided genetic evidence for the relevance of each isoform of the HIF and PHD gene families for oxygen-dependent Epo gene expression.

Two independent inducible PHD2 knockout mouse models showed increased erythropoiesis and angiogenesis as well as evidence for dilated cardiomyopathy and premature death (90, 132, 133). These findings are in line with in vitro data demonstrating that PHD2 is the main oxygen sensor responsible for normoxic HIF- α turnover (16). Consistently, systemic PHD2 but not PHD1 or PHD3 knockout mice die during embryonic development (134). A further confirmation for the major role PHD2 is playing in oxygen-regulated Epo synthesis in vivo comes from the identification of gene mutations that cause familial erythrocytosis: the affected amino acids identified to date confer residues P317, R371, and H374 of PHD2 (71, 103, 106). Neither PHD1 nor PHD3 mutations in familial erythrocytosis have been reported so far.

Due to its nature as a tumor suppressor, a large number of mutations in the human *VHL* gene were known before its role as a HIF- α E3 ubiquitin ligase was discovered. In accordance with its essential and nonredundant role in normoxic HIF- α degradation, it did not come as a surprise that such mutations also provoke erythrocytosis (145). Interestingly, Chuvash polycythemia is caused by a VHL R200W mutation in the absence of cancer (3). A number of additional VHL mutations causing congenital erythrocytosis have been identified thereafter (12, 21, 101, 108). Finally, a genetically altered mouse model recapitulated the polycythemic phenotype (45).

The first evidence that HIF-2 α rather than HIF-1 α might regulate renal Epo synthesis came from descriptive studies by Rosenberg and coworkers (115), who convincingly demonstrated in situ overlapping of HIF-2 α but not HIF-1 α with the peritubular, interstitial REPOS cells. Similar results have recently been reported using chemical PHD inhibition (99). The critical role of HIF-2 α was further confirmed by RNAi experiments in liver-derived hepatoma (140). Because it is not yet possible to target the REPOS cells in mice and since HIF-1 α and HIF-2 α knockout mice do not survive embryonic development (55, 135), knockout mouse models initially provided only circumstantial evidence for the role of each HIF- α isoform in Epo gene expression. Using liver-specific mouse gene targeting, VHL deficiency has been shown to cause polycythemia. This phenotype could be reverted by a VHL-HIF- β /

Fig. 1. Simplified schematic overview of the oxygen signaling cascade in renal Epo-producing and oxygen sensing (REPOS) cells. A: renal tubular epithelial oxygen consumption and countercurrent exchange of oxygen leads to a stable corticomedullary oxygen gradient (blue background). Under hypoxic conditions, this gradient is shifted to the cortex, leading to increased recruitment of REPOS cells and elevated Epo production. B: oxygen partial pressure as well as Krebs cycle intermediates, reactive oxygen species (ROS), and a large number of interacting proteins regulate the abundance and activity of the oxygen-sensing PHD/FIH protein hydroxylases as discussed in detail in the text.

ARNT but not by a VHL-HIF-1 α double knockout, suggesting that HIF-2 α is responsible for Epo induction, at least in the liver (113). These results were subsequently confirmed in a hepatic HIF-2 α knockout mouse model (112). Quite surprisingly, an independent HIF-2 α global knockout mouse strain turned out to be viable, and these mice were indeed anemic with decreased renal Epo expression (124). Finally, conditional global deletion of HIF-2 α but not HIF-1 α resulted in anemia in adult mice (43). The mouse data have recently been confirmed by the identification of HIF-2 α P534L, M535V, M535I, G537W, G537R, and D539E mutations that cause human familial erythrocytosis (36, 37, 80, 102, 104, 107, 138). No HIF-1 α or HIF-3 α mutations have been reported so far to be involved in erythrocytosis.

Altogether, there is now compelling evidence that the oxygen-signaling pathway regulating renal Epo synthesis consists of the PHD2-VHL-HIF-2 α axis. Recent RNAi-based studies confirmed the major role of PHD2 in Epo regulation in vitro as well as in vivo and showed that FIH and the other PHD family members can play a modulatory role in Epo gene expression (32, 98). However, to directly investigate the function of these proteins in the kidney, a transgenic mouse model would be required which targets specifically the REPOS cells.

Multimodal Regulation of Oxygen Sensing

Adding more complexity to renal oxygen sensing, the ratio between PHD and HIF- α levels is interconnected. If either one is upregulated, it overcomes the function of the other. As well as an increase in PHD synthesis that leads to HIF- α degradation, an increase in HIF- α synthesis leads to its own stabilization by saturating the degradation machinery. PHDs have a low oxygen affinity, with K_m values that correspond roughly to the P_{O_2} of room air, which is clearly higher than the highest renal tissue P_{O_2} (46). Thus even under hyperoxic conditions the PHDs are not fully active and a further decrease in the oxygenation as well as a decrease in the PHD levels allow for the stabilization of HIF- α (41, 64, 91, 131). This feature explains why regulation of the PHD levels likely is of physiological relevance: even the slightest up- or downregulation of PHD abundance and/or activity potentially affects HIF-2-dependent Epo expression. Indeed, PHDs are regulated on three levels: transcription, protein abundance/stability, and enzymatic activity.

PHD2 and PHD3, but not PHD1 or FIH, are HIF target genes induced under hypoxic conditions (5, 16, 23, 26, 27, 31, 81, 88, 109). To date, little is known about the tissue-specific and signal-related expression of the genes encoding PHDs: estradiol and LIF are known to induce PHD1 and PHD2 gene expression, respectively, and transforming growth factor (TGF)- β 1 has been shown to inhibit PHD2 gene expression (4, 18, 87, 127).

In addition to transcriptional regulation, the members of the oxygen-signaling cascade can also be regulated at the posttranscriptional level. An iron response element has been identified in the 5'-untranslated region of HIF-2 α mRNA. Iron-regulatory proteins bind to this element and inhibit HIF-2 α translation, thereby suppressing renal Epo production when iron deficiency limits efficient hematopoiesis (120). The Epo mRNA is hypoxically stabilized, contributing to its enormous hypoxic inducibility (42). Epo mRNA binding proteins have

been identified, and the binding site has been shown to be required for hypoxic inducibility (86, 114, 121). More recently, microRNAs (miRNAs) emerged as regulators of the oxygen-signaling cascade: the 3'-untranslated regions of FIH, VHL, and HIF-1 α mRNA are targeted by miR-31, miR-92-1, and miR-20b/miR-199a, respectively (40, 73, 74, 111). However, no miRNAs targeting PHD2, HIF-2 α , or Epo have been reported so far, and how miRNAs are involved in REPOS cell-specific oxygen sensing and Epo expression remains to be investigated.

A number of proteins have recently been identified to interact with PHDs (Table 1). The newly identified interaction partners are able to regulate various aspects of PHD function, including stability, folding, subcellular localization, and enzymatic activity (142). One group of interaction partners serves as a molecular scaffold and includes osteosarcoma amplified 9 (OS-9), A-kinase anchor proteins (AKAP 6 and 12), mitogen-activated protein kinase organizer 1 (Morg1), inhibitor of growth (ING4), and melanoma antigens (MAGE 9 and 11) (6, 8, 22, 24, 44, 48, 97, 147). Other interacting proteins regulate PHD protein levels, including the RING finger E3 ubiquitin ligase seven in absentia homolog 2 (Siah2), the FK506-binding protein 38 (FKBP38), and the chaperonin TCP-1 ring complex (TRiC) (10, 11, 82, 92, 93, 136). Importin- α 5 and CRM1 are involved in PHD1 nuclear import and PHD2 nuclear export, respectively (130). The functional importance of the nuclear-cytoplasmic distribution of PHD2 has recently been demonstrated for the growth of cancer cells (60).

The Siah1 family member additionally regulates FIH protein levels, suggesting that Siah proteins also play a role in fine-tuning the transcriptional activity of HIFs (34, 35). FIH activity is further suppressed by binding to neuronal munc18-1-interacting protein 3 (Mint3)/amyloid β (A4), precursor protein-

Table 1. Putative functions of proteins binding to PHD and FIH oxygen sensors

Interactor	Target	Putative Function	Reference(s)
Protein stability			
Siah1a/2	PHD1/3	PHD degradation	93
Siah1	FIH	FIH degradation	35
FKBP38	PHD2	PHD degradation	10, 11
TRiC	PHD3	Chaperonin	82
Scaffolding proteins			
mAKAP12	PHD2/3	Molecular scaffold	22, 147
ING4	PHD2	HIF- α inhibition	24, 97
OS-9	PHD2/3	Molecular scaffold	8
MAGE-9/11	PHD2	PHD inhibition	6
Morg1	PHD3	Molecular scaffold	44, 48
Importin- α 5	PHD1	Nuclear import	130
CRM1	PHD2	Nuclear export	130
Mint3/ABPA3	FIH	FIH inhibition	119
Others			
VHL	FIH	HIF- α inhibition	76
Cdr2	PHD1	HIF- α inhibition	9
IOP1	PHD2	Induces HIF-1 α mRNA	50

PHD, prolyl-4-hydroxylase domain; FIH, factor-inhibiting HIF; HIF- α , hypoxia-inducible transcription factor- α ; Siah, seven in absentia homolog; FKBP38, FK506-binding protein 38; TRiC, TCP-1 ring complex; AKAP, A-kinase anchor protein; ING, inhibitor of growth; OS-9, osteosarcoma amplified 9; MAGE, melanoma antigen; Morg1, mitogen-activated protein kinase organizer 1; Mint3, munc18-1-interacting protein 3; Cdr2, cerebellar degeneration-related protein 2; IOP1, iron-only hydrogenase-like protein 1; VHL, von Hippel-Lindau.

binding family A member 3 (APBA3), leading to increased HIF activity (119). Intriguingly, FIH has been reported to recruit VHL to HIF, which inhibits HIF transcriptional activity (76). Of note, also PHD2 appears to directly inhibit HIF-1 transcriptional activity, independently of protein hydroxylation and without affecting HIF-1 α protein levels (128, 137). Two other PHD interactors, the iron-only hydrogenase-like protein 1 (IOP1) and the onconeural cerebellar degeneration-related protein 2 (Cdr2) additionally affect HIF- α mRNA and protein, respectively, by unknown mechanisms (9, 50).

PHD-dependent protein hydroxylation is a multicomponent reaction, allowing for the integration of further signaling pathways (94). Indeed, it has been shown that small molecules, such as ascorbate, transition metals, and ROS, including nitric oxide, affect PHD activity, establishing molecular cross talk between oxygen homeostasis and redox-active substances (13, 39, 47, 65, 79, 89). Of major physiological interest is the finding that Krebs cycle intermediates interfere with PHD function (25, 75, 125). Germline mutations of the genes encoding fumarate hydratase (FH) or succinate dehydrogenase (SDH) result in the accumulation of fumarate and succinate, respectively, two potent inhibitors of PHD activity (53, 110). Moreover, mutations in the gene encoding isocitrate dehydrogenase-1 (IDH-1) lead to a decrease in the PHD cosubstrate 2-oxoglutarate and an increase in HIF-1 α levels (148). Thus the PHDs link oxygen sensing with mitochondrial metabolism.

Intrinsic Feedback Mechanisms Trigger Hematocrit-Controlled Renal Epo Synthesis

Figure 1B schematically summarizes the cross talk between the oxygen-sensing and other signaling pathways as mentioned above. However, the physiological relevance of these findings for oxygen-regulated renal Epo expression is currently unknown and awaits appropriate kidney-derived cell culture and mouse models. Since the majority of hereditary erythrocytosis cases are idiopathic diseases (105), this overview might provide the basis for screening additional gene mutations in these patients. Conversely, the identification of such mutations would unequivocally demonstrate the physiological relevance of these genes in renal oxygen sensing.

Regarding the HIF-PHD regulatory loop, it is of interest that hypoxically induced serum Epo levels decrease clearly before the hematocrit is increased, excluding an early negative feedback loop via the subsequent increase in blood oxygen content (1). The drop in serum Epo is paralleled by a drop in renal Epo protein and mRNA levels and is independent of changes in extracellular Epo levels (29, 57). These data suggest one or several intrinsic negative feedback loop(s) that control renal Epo synthesis. A very likely mechanism, albeit unproven regarding Epo regulation, might consist of the HIF-dependent transcriptional increase in PHD2 and PHD3 levels, which in turn downregulate HIF- α protein levels (Fig. 1B). This model shows a biphasic response, with an early PHD3 and a late PHD2 induction, and it allows for ongoing oxygen sensing, suggesting that it could easily adapt to both acute and chronic changes in oxygenation (41, 64, 91, 131).

Neuronal Phenotype of REPOS Cells?

The lack of an appropriate cell culture model seriously hinders further investigation into the physiological relevance of

the complex oxygen-signaling (sub)pathways, as outlined above, for regulation of renal Epo synthesis. Most of our knowledge is based on findings derived from the human hepatoma cell lines HepG2 and Hep3B. Obviously, caution is required when one is transferring these findings to REPOS cells. So far, it was not possible to establish a kidney-derived cell culture system capable of oxygen-regulated Epo synthesis *in vitro*. Extensive *in situ* analyses of the REPOS cells suggested a fibroblast-like phenotype (7, 84). However, fibroblasts would not be considered to be impossible to cultivate. Moreover, despite the many reports on extrarenal Epo expression, fibroblasts have not been shown to express Epo in any other organ than the kidney. Currently, the evidence for a fibroblast-like phenotype comes from renal coexpression of Epo and HIF-2 α together with CD73/ecto-5'-nucleotidase that is considered to be a fibroblast marker (7, 84, 99). However, even under severely hypoxic conditions only a subset of the CD73/ecto-5'-nucleotidase-positive cells is also positive for HIF-2 α /Epo, and CD73/ecto-5'-nucleotidase is not solely expressed by fibroblasts (according to the Human Protein Atlas, <http://www.proteinatlas.org>; and the GNF atlas, <http://biogps.gnf.org>), suggesting that REPOS cells could also be of a different nature.

Intriguingly, a recent study using genetically modified mice that carry a green fluorescent protein (GFP) reporter protein under the control of a 180-kb Epo transgene suggested a "neuron-like" phenotype (95). The GFP-positive cells have dendrite-like processes and express the neuronal markers microtubule-associated protein 2 (MAP2) and neurofilament light polypeptide. A neuron-like phenotype would explain the failure of *in vitro* cultivation of these cells and is in line with their sensory function. Remarkably, in the central nervous system neurons and astrocytes are known to express Epo (14, 78, 83).

There is also an interesting analogy between chemosensory cells of the kidney and carotid body. Both oxygen-signaling pathways need the more infrequently expressed HIF-2 α isoform. The dopaminergic type I cells of the carotid body rely on HIF-2 α for hypoxia-inducible expression of tyrosine hydroxylase, the rate-limiting enzyme in catecholamine synthesis: HIF-2 α knockout mice die from bradycardia due to the lack of catecholamines (135); HIF-2 α knockdown results in decreased catecholamine synthesis in adrenomedullary chromaffin cells (19); HIF-2 α , but not HIF-1 α , deficiency reverts the reduced apoptosis and increased cell numbers in the sympathoadrenal system, including the carotid body, of PHD3 knockout mice (17); and mutations in the *VHL* and *SDH* genes are associated with an increased risk of paraganglioma, including carotid body tumors (61). Considering this analogy, kidney-derived cell culture systems might profit from the recent finding that glia-like stem cells generate neurotropic factors that sustain carotid body neurogenesis (100). All in all, there might be new hope for a REPOS cell culture model.

Tissue-Specific Gene Expression in REPOS Cells

Another important question that is difficult to address due to the lack of a REPOS cell culture model is the tissue specificity of oxygen-sensing and -signaling factors in REPOS cells. Whereas PHD2 and VHL are ubiquitously expressed, HIF-2 α and Epo show highly REPOS cell-specific expression patterns on the mRNA level. As mentioned above, renal oxygen sensing

and signaling, mandatory to systemic red blood cell and oxygen homeostasis, depends on the correct spatial localization of the cells expressing Epo within the renal oxygen gradient. REPOS cell-restricted gene expression is equally important to proper oxygen sensing as hypoxia inducibility itself. While virtually nothing is known about the REPOS cell-specific gene expression of HIF-2 α , a number of transcription factors have been implicated in the tissue-specific expression of the Epo gene (28). Whether similar mechanisms are involved in HIF-2 α expression in REPOS cells remains to be investigated.

Under normoxic conditions, Epo gene expression is strikingly low, consistent with the idea that the conditional (i.e., hypoxic) HIF-2-mediated induction of the Epo gene plays a major role in its REPOS cell-specific expression. Epigenetic modifications by CpG DNA methylation are likely to be involved in the constitutive suppression of Epo expression in non-REPOS cell types. Of note, it has been shown that HIF binding to its consensus DNA recognition site in the Epo 3'-HRE is CpG DNA methylation sensitive (117, 143). At least for hepatocytes, HIF-2 cooperation with hepatic nuclear factor 4 (HNF-4) at the 3' HRE might contribute to tissue-specific gene expression (38). However, how HNF-4 contributes to renal Epo expression is less clear, and it should be remembered that the importance of the Epo 3' HRE for REPOS cell-specific gene expression awaits a formal proof. Very low basal Epo expression also suggests the involvement of transcriptional suppressors. Indeed, GATA factors have been shown to repress Epo gene promoter activity (51, 52). Using transgenic mice, it has been demonstrated that the GATA *cis*-regulatory element of the Epo promoter is important for repression of Epo gene expression in non-REPOS cells of the kidney (95).

Conclusions

Albeit not fully understood regarding renal Epo production, the wealth of novel insights into the basic mechanisms of oxygen sensing and signaling opened new routes to the treatment of diseases related to oxygen sensing. In kidney failure, oxygen consumption is decreased, resulting in increased tissue PO₂, decreased renal Epo production, and anemia. Kidney dialysis patients hence need lifelong Epo treatments. Pharmaceutical interference with PHD/FIH enzymatic activity proved to be efficient *in vitro* and beneficial for the treatment of a number of diseases in animal models *in vivo*, including tissue ischemia, kidney injury, and transplantation (15, 33, 63, 99, 129). Protein hydroxylase inhibitors also stimulated Epo production in mouse and rhesus macaque models (49, 118). Clinical phase II studies are being conducted to test the safety and efficacy of protein hydroxylase inhibitors in the treatment of patients with renal anemia (ClinicalTrials.gov identifiers NCT00456053 and NCT00761657). Even if it is currently unclear to what extent Epo is derived from the liver and to what extent from the kidney following treatment with these protein hydroxylase inhibitors, these examples demonstrate the clinical relevance of basic research in renal oxygen sensing. Studies in the near future will answer the question as to whether the currently available inhibitors of the oxygen-sensing protein hydroxylases are sufficiently efficient and specific to serve as novel therapy strategies for the treatment of anemia caused by renal disease.

GRANTS

The authors' work is supported by the Swiss National Science Foundation (Grant 3100AO-116047/1) and the Sassella Stiftung. D. Hoogewijs is a recipient of a Marie Curie Fellowship from the European Commission.

DISCLOSURES

No conflicts of interest are declared by the authors.

REFERENCES

1. **Abbrecht PH, Littell JK.** Plasma erythropoietin in men and mice during acclimatization to different altitudes. *J Appl Physiol* 32: 54–58, 1972.
2. **Acker T, Fandrey J, Acker H.** The good, the bad and the ugly in oxygen-sensing: ROS, cytochromes and prolyl-hydroxylases. *Cardiovasc Res* 71: 195–207, 2006.
3. **Ang SO, Chen H, Hirota K, Gordeuk VR, Jelinek J, Guan Y, Liu E, Sergueeva AI, Miasnikova GY, Mole D, Maxwell PH, Stockton DW, Semenza GL, Prchal JT.** Disruption of oxygen homeostasis underlies congenital Chuvash polycythemia. *Nat Genet* 32: 614–621, 2002.
4. **Appelhoff RJ, Tian YM, Raval RR, Turley H, Harris AL, Pugh CW, Ratcliffe PJ, Gleadle JM.** Differential function of the prolyl hydroxylases PHD1, PHD2, and PHD3 in the regulation of hypoxia-inducible factor. *J Biol Chem* 279: 38458–38465, 2004.
5. **Aprelikova O, Chandramouli GV, Wood M, Vasselli JR, Riss J, Maranchie JK, Linehan WM, Barrett JC.** Regulation of HIF prolyl hydroxylases by hypoxia-inducible factors. *J Cell Biochem* 92: 491–501, 2004.
6. **Aprelikova O, Pandolfi S, Tackett S, Ferreira M, Salnikow K, Ward Y, Risinger JJ, Barrett JC, Niederhuber J.** Melanoma antigen-11 inhibits the hypoxia-inducible factor prolyl hydroxylase 2 and activates hypoxic response. *Cancer Res* 69: 616–624, 2009.
7. **Bachmann S, Le Hir M, Eckardt KU.** Co-localization of erythropoietin mRNA and ecto-5'-nucleotidase immunoreactivity in peritubular cells of rat renal cortex indicates that fibroblasts produce erythropoietin. *J Histochem Cytochem* 41: 335–341, 1993.
8. **Baek JH, Mahon PC, Oh J, Kelly B, Krishnamachary B, Pearson M, Chan DA, Giaccia AJ, Semenza GL.** OS-9 interacts with hypoxia-inducible factor 1 α and prolyl hydroxylases to promote oxygen-dependent degradation of HIF-1 α . *Mol Cell* 17: 503–512, 2005.
9. **Balamurugan K, Luu VD, Kaufmann MR, Hofmann VS, Boysen G, Barth S, Bordoli MR, Stiehl DP, Moch H, Schraml P, Wenger RH, Camenisch G.** Onconeuronal cerebellar degeneration-related antigen, Cdr2, is strongly expressed in papillary renal cell carcinoma and leads to attenuated hypoxic response. *Oncogene* 28: 3274–3285, 2009.
10. **Barth S, Edlich F, Berchner-Pfannschmidt U, Gneuss S, Jahreis G, Hasgall PA, Fandrey J, Wenger RH, Camenisch G.** Hypoxia-inducible factor prolyl-4-hydroxylase PHD2 protein abundance depends on iIntegral membrane anchoring of FKBP38. *J Biol Chem* 284: 23046–23058, 2009.
11. **Barth S, Nesper J, Hasgall PA, Wirthner R, Nytko KJ, Edlich F, Katschinski DM, Stiehl DP, Wenger RH, Camenisch G.** The peptidyl prolyl *cis/trans* isomerase FKBP38 determines hypoxia-inducible transcription factor prolyl-4-hydroxylase PHD2 protein stability. *Mol Cell Biol* 27: 3758–3768, 2007.
12. **Bento MC, Chang KT, Guan Y, Liu E, Caldas G, Gatti RA, Prchal JT.** Congenital polycythemia with homozygous and heterozygous mutations of von Hippel-Lindau gene: five new Caucasian patients. *Haematologica* 90: 128–129, 2005.
13. **Berchner-Pfannschmidt U, Yamac H, Trinidad B, Fandrey J.** Nitric oxide modulates oxygen sensing by hypoxia-inducible factor 1-dependent induction of prolyl hydroxylase 2. *J Biol Chem* 282: 1788–1796, 2007.
14. **Bernaudo M, Bellail A, Marti HH, Yvon A, Vivien D, Duchatelle I, Mackenzie ET, Petit E.** Neurons and astrocytes express EPO mRNA: oxygen-sensing mechanisms that involve the redox-state of the brain. *Glia* 30: 271–278, 2000.
15. **Bernhardt WM, Gottmann U, Doyon F, Buchholz B, Campean V, Schödel J, Reisenbuechler A, Klaus S, Arend M, Flippin L, Willam C, Wiesener MS, Yard B, Warnecke C, Eckardt KU.** Donor treatment with a PHD-inhibitor activating HIFs prevents graft injury and prolongs survival in an allogenic kidney transplant model. *Proc Natl Acad Sci USA* 106: 21276–21281, 2009.

16. Berra E, Benizri E, Ginouves A, Volmat V, Pouyssegur J. HIF prolyl-hydroxylase 2 is the key oxygen sensor setting low steady-state levels of HIF-1 α in normoxia. *EMBO J* 22: 4082–4090, 2003.
17. Bishop T, Gallagher D, Pascual A, Lygate CA, de Bono JP, Nicholls LG, Ortega-Saenz P, Oster H, Wijeyekoon B, Sutherland AI, Grosfeld A, Aragonés J, Schneider M, van Geyte K, Teixeira D, Diez-Juan A, Lopez-Barneo J, Channon KM, Maxwell PH, Pugh CW, Davies AM, Carmeliet P, Ratcliffe PJ. Abnormal sympathoadrenal development and systemic hypotension in PHD3^{-/-} mice. *Mol Cell Biol* 28: 3386–3400, 2008.
18. Bozec A, Bakiri L, Hoebertz A, Eferl R, Schilling AF, Komnenovic V, Scheuch H, Priemel M, Stewart CL, Amling M, Wagner EF. Osteoclast size is controlled by Fra-2 through LIF/LIF-receptor signaling and hypoxia. *Nature* 454: 221–225, 2008.
19. Brown ST, Kelly KF, Daniel JM, Nurse CA. Hypoxia inducible factor (HIF)-2 α is required for the development of the catecholaminergic phenotype of sympathoadrenal cells. *J Neurochem* 110: 622–630, 2009.
20. Bruick RK, McKnight SL. A conserved family of prolyl-4-hydroxylases that modify HIF. *Science* 294: 1337–1340, 2001.
21. Cario H, Schwarz K, Jorch N, Kyank U, Petrides PE, Schneider DT, Uhle R, Debatin KM, Kohne E. Mutations in the von Hippel-Lindau (VHL) tumor suppressor gene and VHL-haplotype analysis in patients with presumable congenital erythrocytosis. *Haematologica* 90: 19–24, 2005.
22. Choi YK, Kim JH, Kim WJ, Lee HY, Park JA, Lee SW, Yoon DK, Kim HH, Chung H, Yu YS, Kim KW. AKAP12 regulates human blood-retinal barrier formation by downregulation of hypoxia-inducible factor-1 α . *J Neurosci* 27: 4472–4481, 2007.
23. Cioffi CL, Liu XQ, Kosinski PA, Garay M, Bowen BR. Differential regulation of HIF-1 α prolyl-4-hydroxylase genes by hypoxia in human cardiovascular cells. *Biochem Biophys Res Commun* 303: 947–953, 2003.
24. Colla S, Tagliaferri S, Morandi F, Lunghi P, Donofrio G, Martorana D, Mancini C, Lazzaretti M, Mazzeri L, Ravanetti L, Bonomini S, Ferrari L, Miranda C, Ladetto M, Neri A, Greco A, Mangoni M, Bonati A, Rizzoli V, Giuliani N. The new tumor-suppressor gene inhibitor of growth family member 4 (ING4) regulates the production of proangiogenic molecules by myeloma cells and suppresses hypoxia-inducible factor-1 α (HIF-1 α) activity: involvement in myeloma-induced angiogenesis. *Blood* 110: 4464–4475, 2007.
25. Dalgard CL, Lu H, Mohyeldin A, Verma A. Endogenous 2-oxoacids differentially regulate expression of oxygen sensors. *Biochem J* 380: 419–424, 2004.
26. D'Angelo G, Duplan E, Boyer N, Vigne P, Frelin C. Hypoxia up-regulates prolyl hydroxylase activity: a feedback mechanism that limits HIF-1 responses during reoxygenation. *J Biol Chem* 278: 38183–38187, 2003.
27. del Peso L, Castellanos MC, Temes E, Martin-Puig S, Cuevas Y, Olmos G, Landázuri MO. The von Hippel Lindau/hypoxia-inducible factor (HIF) pathway regulates the transcription of the HIF-proline hydroxylase genes in response to low oxygen. *J Biol Chem* 278: 48690–48695, 2003.
28. Ebert BL, Bunn HF. Regulation of the erythropoietin gene. *Blood* 94: 1864–1877, 1999.
29. Eckardt KU, Dittmer J, Neumann R, Bauer C, Kurtz A. Decline of erythropoietin formation at continuous hypoxia is not due to feedback inhibition. *Am J Physiol Renal Physiol* 278: F1432–F1437, 1999.
30. Eckardt KU, Koury ST, Tan CC, Schuster SJ, Kaissling B, Ratcliffe PJ, Kurtz A. Distribution of erythropoietin producing cells in rat kidneys during hypoxic hypoxia. *Kidney Int* 43: 815–823, 1993.
31. Epstein AC, Gleadle JM, McNeill LA, Hewitson KS, O'Rourke J, Mole DR, Mukherji M, Metzén E, Wilson MI, Dhanda A, Tian YM, Masson N, Hamilton DL, Jaakkola P, Barstead R, Hodgkin J, Maxwell PH, Pugh CW, Schofield CJ, Ratcliffe PJ. *C. elegans* EGL-9 and mammalian homologs define a family of dioxygenases that regulate HIF by prolyl hydroxylation. *Cell* 107: 43–54, 2001.
32. Fisher TS, Lira PD, Stock JL, Perregaux DG, Brissette WH, Ozolins TR, Li B. Analysis of the role of the HIF hydroxylase family members in erythropoiesis. *Biochem Biophys Res Commun* 388: 683–688, 2009.
33. Fraisl P, Aragonés J, Carmeliet P. Inhibition of oxygen sensors as a therapeutic strategy for ischaemic and inflammatory disease. *Nat Rev Drug Discov* 8: 139–152, 2009.
34. Fukuba H, Takahashi T, Jin HG, Kohriyama T, Matsumoto M. Abundance of asparaginyl-hydroxylase FIH is regulated by Siah-1 under normoxic conditions. *Neurosci Lett* 433: 209–214, 2008.
35. Fukuba H, Yamashita H, Nagano Y, Jin HG, Hiji M, Ohtsuki T, Takahashi T, Kohriyama T, Matsumoto M. Siah-1 facilitates ubiquitination and degradation of factor inhibiting HIF-1 α (FIH). *Biochem Biophys Res Commun* 353: 324–329, 2007.
36. Furlow PW, Percy MJ, Sutherland S, Bierl C, McMullin MF, Master SR, Lappin TR, Lee FS. Erythrocytosis-associated HIF-2 α mutations demonstrate a critical role for residues C-terminal to the hydroxylacceptor proline. *J Biol Chem* 284: 9050–9058, 2009.
37. Gale DP, Harten SK, Reid CD, Tuddenham EG, Maxwell PH. Autosomal dominant erythrocytosis and pulmonary arterial hypertension associated with an activating HIF2 α mutation. *Blood* 112: 919–921, 2008.
38. Galson DL, Tsuchiya T, Tendler DS, Huang LE, Ren Y, Ogura T, Bunn HF. The orphan receptor hepatic nuclear factor 4 functions as a transcriptional activator for tissue-specific and hypoxia-specific erythropoietin gene expression and is antagonized by EAR3/COUP-TF1. *Mol Cell Biol* 15: 2135–2144, 1995.
39. Gerald D, Berra E, Frapart YM, Chan DA, Giaccia AJ, Mansuy D, Pouyssegur J, Yaniv M, Mechta-Grigoriou F. JunD reduces tumor angiogenesis by protecting cells from oxidative stress. *Cell* 118: 781–794, 2004.
40. Ghosh AK, Shanafelt TD, Cimmino A, Taccioli C, Volinia S, Liu CG, Calin GA, Croce CM, Chan DA, Giaccia AJ, Secretò C, Wellik LE, Lee YK, Mukhopadhyay D, Kay NE. Aberrant regulation of pVHL levels by microRNA promotes the HIF/VEGF axis in CLL B cells. *Blood* 113: 5568–5574, 2009.
41. Ginouves A, Ilc K, Macias N, Pouyssegur J, Berra E. PHDs overactivation during chronic hypoxia “desensitizes” HIF α and protects cells from necrosis. *Proc Natl Acad Sci USA* 105: 4745–4750, 2008.
42. Goldberg MA, Gaut CC, Bunn HF. Erythropoietin mRNA levels are governed by both the rate of gene transcription and posttranscriptional events. *Blood* 77: 271–277, 1991.
43. Gruber M, Hu CJ, Johnson RS, Brown EJ, Keith B, Simon MC. Acute postnatal ablation of Hif-2 α results in anemia. *Proc Natl Acad Sci USA* 104: 2301–2306, 2007.
44. Hammerschmidt E, Loeffler I, Wolf G. Morgl heterozygous mice are protected from acute renal ischemia-reperfusion injury. *Am J Physiol Renal Physiol* 297: F1273–F1287, 2009.
45. Hickey MM, Lam JC, Bezman NA, Rathmell WK, Simon MC. von Hippel-Lindau mutation in mice recapitulates Chuvash polycythemia via hypoxia-inducible factor-2 α signaling and splenic erythropoiesis. *J Clin Invest* 117: 3879–3889, 2007.
46. Hirsilä M, Koivunen P, Günzler V, Kivirikko KI, Myllyharju J. Characterization of the human prolyl 4-hydroxylases that modify the hypoxia-inducible factor. *J Biol Chem* 278: 30772–30780, 2003.
47. Hirsilä M, Koivunen P, Xu L, Seeley T, Kivirikko KI, Myllyharju J. Effect of desferrioxamine and metals on the hydroxylases in the oxygen sensing pathway. *FASEB J* 19: 1308–1310, 2005.
48. Hopfer U, Hopfer H, Jablonski K, Stahl RA, Wolf G. The novel WD-repeat protein Morgl acts as a molecular scaffold for hypoxia-inducible factor prolyl hydroxylase 3 (PHD3). *J Biol Chem* 281: 8645–8655, 2006.
49. Hsieh MM, Linde NS, Wynter A, Metzger M, Wong C, Langsetmo I, Lin A, Smith R, Rodgers GP, Donahue RE, Klaus SJ, Tisdale JF. HIF prolyl hydroxylase inhibition results in endogenous erythropoietin induction, erythrocytosis, and modest fetal hemoglobin expression in rhesus macaques. *Blood* 110: 2140–2147, 2007.
50. Huang J, Song D, Flores A, Zhao Q, Mooney SM, Shaw LM, Lee FS. IOPI, a novel hydrogenase-like protein that modulates hypoxia-inducible factor-1 α activity. *Biochem J* 401: 341–352, 2007.
51. Imagawa S, Suzuki N, Ohmine K, Obara N, Mukai HY, Ozawa K, Yamamoto M, Nagasawa T. GATA suppresses erythropoietin gene expression through GATA site in mouse erythropoietin gene promoter. *Int J Hematol* 75: 376–381, 2002.
52. Imagawa S, Yamamoto M, Miura Y. Negative regulation of the erythropoietin gene expression by the GATA transcription factors. *Blood* 89: 1430–1439, 1997.
53. Isaacs JS, Jung YJ, Mole DR, Lee S, Torres-Cabala C, Chung YL, Merino M, Trepel J, Zbar B, Toro J, Ratcliffe PJ, Linehan WM, Neckers L. HIF overexpression correlates with biallelic loss of fumarate

- hydratase in renal cancer: novel role of fumarate in regulation of HIF stability. *Cancer Cell* 8: 143–153, 2005.
54. Ivan M, Kondo K, Yang H, Kim W, Valiando J, Ohh M, Salic A, Asara JM, Lane WS, Kaelin WG Jr. HIF α targeted for VHL-mediated destruction by proline hydroxylation: implications for O₂ sensing. *Science* 292: 464–468, 2001.
 55. Iyer NV, Kotch LE, Agani F, Leung SW, Laughner E, Wenger RH, Gassmann M, Gearhart JD, Lawler AM, Yu AY, Semenza GL. Cellular and developmental control of O₂ homeostasis by hypoxia-inducible factor 1 α . *Genes Dev* 12: 149–162, 1998.
 56. Jaakkola P, Mole DR, Tian YM, Wilson MI, Gielbert J, Gaskell SJ, von Kriegsheim A, Hebestreit HF, Mukherji M, Schofield CJ, Maxwell PH, Pugh CW, Ratcliffe PJ. Targeting of HIF- α to the von Hippel-Lindau ubiquitylation complex by O₂-regulated prolyl hydroxylation. *Science* 292: 468–472, 2001.
 57. Jelkmann W. Temporal pattern of erythropoietin titers in kidney tissue during hypoxic hypoxia. *Eur J Physiol* 393: 88–91, 1982.
 58. Jelkmann W. Erythropoietin: structure, control of production, and function. *Physiol Rev* 72: 449–489, 1992.
 59. Jewell UR, Kvietikova I, Scheid A, Bauer C, Wenger RH, Gassmann M. Induction of HIF-1 α in response to hypoxia is instantaneous. *FASEB J* 15: 1312–1314, 2001.
 60. Jokilehto T, Högel H, Heikkinen P, Rantanen K, Elenius K, Sundström J, Jaakkola PM. Retention of prolyl hydroxylase PHD2 in the cytoplasm prevents PHD2-induced anchorage-independent carcinoma cell growth. *Exp Cell Res* 316: 1169–1178, 2010.
 61. Kaelin WG. Von Hippel-Lindau disease. *Annu Rev Pathol* 2: 145–173, 2007.
 62. Kaelin WG Jr, Ratcliffe PJ. Oxygen sensing by metazoans: the central role of the HIF hydroxylase pathway. *Mol Cell* 30: 393–402, 2008.
 63. Katschinski DM. In vivo functions of the prolyl-4-hydroxylase domain oxygen sensors: direct route to the treatment of anaemia and the protection of ischaemic tissues. *Acta Physiol (Oxf)* 195: 407–414, 2009.
 64. Khanna S, Roy S, Maurer M, Ratan RR, Sen CK. Oxygen-sensitive reset of hypoxia-inducible factor transactivation response: prolyl hydroxylases tune the biological normoxic set point. *Free Radic Biol Med* 40: 2147–2154, 2006.
 65. Knowles HJ, Raval RR, Harris AL, Ratcliffe PJ. Effect of ascorbate on the activity of hypoxia-inducible factor in cancer cells. *Cancer Res* 63: 1764–1768, 2003.
 66. Koivunen P, Tiainen P, Hyvarinen J, Williams KE, Sormunen R, Klaus SJ, Kivirikko KI, Myllyharju J. An endoplasmic reticulum transmembrane prolyl 4-hydroxylase is induced by hypoxia and acts on hypoxia-inducible factor α . *J Biol Chem* 282: 30544–30552, 2007.
 67. Koury MJ, Bondurant MC, Graber SE, Sawyer ST. Erythropoietin messenger RNA levels in developing mice and transfer of ¹²⁵I-erythropoietin by the placenta. *J Clin Invest* 82: 154–159, 1988.
 68. Koury ST, Bondurant MC, Koury MJ. Localization of erythropoietin synthesizing cells in murine kidneys by in situ hybridization. *Blood* 71: 524–527, 1988.
 69. Koury ST, Koury MJ, Bondurant MC, Caro J, Graber SE. Quantitation of erythropoietin-producing cells in kidneys of mice by in situ hybridization: correlation with hematocrit, renal erythropoietin mRNA, and serum erythropoietin concentration. *Blood* 74: 645–651, 1989.
 70. Lacombe C, Da Silva JL, Bruneval P, Fournier JG, Wendling F, Casadevall N, Camilleri JP, Bariety J, Varet B, Tambourin P. Peritubular cells are the site of erythropoietin synthesis in the murine hypoxic kidney. *J Clin Invest* 81: 620–623, 1988.
 71. Ladroue C, Carcenac R, Leporrier J, Gad S, Le Hello C, Galateau-Salle F, Feunteun J, Pouyssegur J, Richard S, Gardie B. PHD2 mutation and congenital erythrocytosis with paraganglioma. *N Engl J Med* 359: 2685–2692, 2008.
 72. Lando D, Peet DJ, Gorman JJ, Whelan DA, Whitelaw ML, Bruick RK. FIH-1 is an asparaginyl hydroxylase enzyme that regulates the transcriptional activity of hypoxia-inducible factor. *Genes Dev* 16: 1466–1471, 2002.
 73. Lei Z, Li B, Yang Z, Fang H, Zhang GM, Feng ZH, Huang B. Regulation of HIF-1 α and VEGF by miR-20b tunes tumor cells to adapt to the alteration of oxygen concentration. *PLoS One* 4: e7629, 2009.
 74. Liu CJ, Tsai MM, Hung PS, Kao SY, Liu TY, Wu KJ, Chiou SH, Lin SC, Chang KW. miR-31 ablates expression of the HIF regulatory factor FIH to activate the HIF pathway in head and neck carcinoma. *Cancer Res* 70: 1635–1644, 2010.
 75. Lu H, Dalgard CL, Mohyeldin A, McFate T, Tait AS, Verma A. Reversible inactivation of HIF-1 prolyl hydroxylases allows cell metabolism to control basal HIF-1. *J Biol Chem* 280: 41928–41939, 2005.
 76. Mahon PC, Hirota K, Semenza GL. FIH-1: a novel protein that interacts with HIF-1 α and VHL to mediate repression of HIF-1 transcriptional activity. *Genes Dev* 15: 2675–2686, 2001.
 77. Makino Y, Kanopka A, Wilson WJ, Tanaka H, Poellinger L. Inhibitory PAS domain protein (IPAS) is a hypoxia-inducible splicing variant of the hypoxia-inducible factor-3 α locus. *J Biol Chem* 277: 32405–32408, 2002.
 78. Marti HH, Wenger RH, Rivas LA, Straumann U, Digicaylioglu M, Henn V, Yonekawa Y, Bauer C, Gassmann M. Erythropoietin gene expression in human, monkey and murine brain. *Eur J Neurosci* 8: 666–676, 1996.
 79. Martin F, Linden T, Katschinski DM, Oehme F, Flamme I, Mukhopadhyay CK, Eckhardt K, Tröger J, Barth S, Camenisch G, Wenger RH. Copper-dependent activation of hypoxia-inducible factor (HIF)-1: implications for ceruloplasmin regulation. *Blood* 105: 4613–4619, 2005.
 80. Martini M, Teofili L, Cenci T, Giona F, Torti L, Rea M, Foa R, Leone G, Larocca LM. A novel heterozygous HIF2A^{M535I} mutation reinforces the role of oxygen sensing pathway disturbances in the pathogenesis of familial erythrocytosis. *Haematologica* 93: 1068–1071, 2008.
 81. Marxsen JH, Stengel P, Doege K, Heikkinen P, Jokilehto T, Wagner T, Jelkmann W, Jaakkola P, Metzén E. Hypoxia-inducible factor-1 (HIF-1) promotes its degradation by induction of HIF- α -prolyl-4-hydroxylases. *Biochem J* 381: 761–767, 2004.
 82. Masson N, Appelhoff RJ, Tuckerman JR, Tian YM, Demol H, Puype M, Vandekerckhove J, Ratcliffe PJ, Pugh CW. The HIF prolyl hydroxylase PHD3 is a potential substrate of the TRiC chaperonin. *FEBS Lett* 570: 166–170, 2004.
 83. Masuda S, Okano M, Yamagishi K, Nagao M, Ueda M, Sasaki R. A novel site of erythropoietin production. Oxygen-dependent production in cultured rat astrocytes. *J Biol Chem* 269: 19488–19493, 1994.
 84. Maxwell PH, Osmond MK, Pugh CW, Heryet A, Nicholls LG, Tan CC, Doe BG, Ferguson DJ, Johnson MH, Ratcliffe PJ. Identification of the renal erythropoietin-producing cells using transgenic mice. *Kidney Int* 44: 1149–1162, 1993.
 85. Maxwell PH, Wiesener MS, Chang GW, Clifford SC, Vaux EC, Cockman ME, Wykoff CC, Pugh CW, Maher ER, Ratcliffe PJ. The tumour suppressor protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis. *Nature* 399: 271–275, 1999.
 86. McGary EC, Rondon IJ, Beckman BS. Post-transcriptional regulation of erythropoietin mRNA stability by erythropoietin mRNA-binding protein. *J Biol Chem* 272: 8628–8634, 1997.
 87. McMahon S, Charbonneau M, Grandmont S, Richard DE, Dubois CM. Transforming growth factor β 1 induces hypoxia-inducible factor-1 stabilization through selective inhibition of PHD2 expression. *J Biol Chem* 281: 24171–24181, 2006.
 88. Metzén E, Stiehl DP, Doege K, Marxsen JH, Hellwig-Bürgel T, Jelkmann W. Regulation of the prolyl hydroxylase domain protein 2 (*phd2/egl-1*) gene: identification of a functional hypoxia-responsive element. *Biochem J* 387: 711–717, 2005.
 89. Metzén E, Zhou J, Jelkmann W, Fandrey J, Brüne B. Nitric oxide impairs normoxic degradation of HIF-1 α by inhibition of prolyl hydroxylases. *Mol Biol Cell* 14: 3470–3481, 2003.
 90. Minamishima YA, Moslehi J, Bardeesy N, Cullen D, Bronson RT, Kaelin WG Jr. Somatic inactivation of the PHD2 prolyl hydroxylase causes polycythemia and congestive heart failure. *Blood* 111: 3236–3244, 2008.
 91. Minamishima YA, Moslehi J, Padera RF, Bronson RT, Liao R, Kaelin WG Jr. A feedback loop involving the Phd3 prolyl hydroxylase tunes the mammalian hypoxic response in vivo. *Mol Cell Biol* 29: 5729–5741, 2009.
 92. Möller A, House CM, Wong CS, Scanlon DB, Liu MC, Ronai Z, Bowtell DD. Inhibition of Siah ubiquitin ligase function. *Oncogene* 28: 289–296, 2009.
 93. Nakayama K, Frew IJ, Hagensen M, Skals M, Habelhah H, Bhoumik A, Kadoya T, Erdjument-Bromage H, Tempst P, Frappell PB, Bowtell DD, Ronai Z. Siah2 regulates stability of prolyl-hydroxylases, controls HIF1 α abundance, and modulates physiological responses to hypoxia. *Cell* 117: 941–952, 2004.

94. Nytko KJ, Spielmann P, Camenisch G, Wenger RH, Stiehl DP. Regulated function of the prolyl-4-hydroxylase domain (PHD) oxygen sensor proteins. *Antioxid Redox Signal* 9: 1329–1338, 2007.
95. Obara N, Suzuki N, Kim K, Nagasawa T, Imagawa S, Yamamoto M. Repression via the GATA box is essential for tissue-specific erythropoietin gene expression. *Blood* 111: 5223–5232, 2008.
96. Oehme F, Ellinghaus P, Kolkhof P, Smith TJ, Ramakrishnan S, Hutter J, Schramm M, Flamme I. Overexpression of PH-4, a novel putative proline 4-hydroxylase, modulates activity of hypoxia-inducible transcription factors. *Biochem Biophys Res Commun* 296: 343–349, 2002.
97. Ozer A, Wu LC, Bruick RK. The candidate tumor suppressor ING4 represses activation of the hypoxia inducible factor (HIF). *Proc Natl Acad Sci USA* 102: 7481–7486, 2005.
98. Ozolins TR, Fisher TS, Nadeau DM, Stock JL, Klein AS, Milici AJ, Morton D, Wilhelms MB, Brissette WH, Li B. Defects in embryonic development of EGLN1/PHD2 knockdown transgenic mice are associated with induction of Igfbp in the placenta. *Biochem Biophys Res Commun* 390: 372–376, 2009.
99. Paliege A, Rosenberger C, Bondke A, Sciesielski L, Shina A, Heyman SN, Flippin LA, Arend M, Klaus SJ, Bachmann S. Hypoxia-inducible factor-2 α -expressing interstitial fibroblasts are the only renal cells that express erythropoietin under hypoxia-inducible factor stabilization. *Kidney Int* 77: 312–318, 2010.
100. Pardal R, Ortega-Sáenz P, Durán R, López-Barneo J. Glia-like stem cells sustain physiologic neurogenesis in the adult mammalian carotid body. *Cell* 131: 364–377, 2007.
101. Pastore YD, Jelinek J, Ang S, Guan Y, Liu E, Jedlickova K, Krishnamurti L, Prchal JT. Mutations in the VHL gene in sporadic apparently congenital polycythemia. *Blood* 101: 1591–1595, 2003.
102. Percy MJ, Beer PA, Campbell G, Dekker AW, Green AR, Oscier D, Rainey MG, van Wijk R, Wood M, Lappin TR, McMullin MF, Lee FS. Novel exon 12 mutations in the HIF2A gene associated with erythrocytosis. *Blood* 111: 5400–5402, 2008.
103. Percy MJ, Furlow PW, Beer PA, Lappin TR, McMullin MF, Lee FS. A novel erythrocytosis-associated PHD2 mutation suggests the location of a HIF binding groove. *Blood* 110: 2193–2196, 2007.
104. Percy MJ, Furlow PW, Lucas GS, Li X, Lappin TR, McMullin MF, Lee FS. A gain-of-function mutation in the HIF2A gene in familial erythrocytosis. *N Engl J Med* 358: 162–168, 2008.
105. Percy MJ, Rumi E. Genetic origins and clinical phenotype of familial and acquired erythrocytosis and thrombocytosis. *Am J Hematol* 84: 46–54, 2009.
106. Percy MJ, Zhao Q, Flores A, Harrison C, Lappin TR, Maxwell PH, McMullin MF, Lee FS. A family with erythrocytosis establishes a role for prolyl hydroxylase domain protein 2 in oxygen homeostasis. *Proc Natl Acad Sci USA* 103: 654–659, 2006.
107. Perrotta S, Della Ragione F. The HIF2A gene in familial erythrocytosis. *N Engl J Med* 358: 1966, 2008.
108. Perrotta S, Nobili B, Ferraro M, Migliaccio C, Borriello A, Cucciolla V, Martinelli V, Rossi F, Punzo F, Cirillo P, Parisi G, Zappia V, Rotoli B, Ragione FD. Von Hippel-Lindau-dependent polycythemia is endemic on the island of Ischia: identification of a novel cluster. *Blood* 107: 514–519, 2006.
109. Pescador N, Cuevas Y, Naranjo S, Alcaide M, Villar D, Landázuri MO, Del Peso L. Identification of a functional hypoxia-responsive element that regulates the expression of the *egl nine* homologue 3 (*egl3/phd3*) gene. *Biochem J* 390: 189–197, 2005.
110. Pollard PJ, Brière JJ, Alam NA, Barwell J, Barclay E, Wortham NC, Hunt T, Mitchell M, Olpin S, Moat SJ, Hargreaves IP, Heales SJ, Chung YL, Griffiths JR, Dalgleish A, McGrath JA, Gleeson MJ, Hodgson SV, Poulson R, Rustin P, Tomlinson IP. Accumulation of Krebs cycle intermediates and over-expression of HIF1 α in tumours which result from germline FH and SDH mutations. *Hum Mol Genet* 14: 2231–2239, 2005.
111. Rane S, He M, Sayed D, Vashistha H, Malhotra A, Sadoshima J, Vatner DE, Vatner SF, Abdellatif M. Downregulation of miR-199a derepresses hypoxia-inducible factor-1 α and Sirtuin 1 and recapitulates hypoxia preconditioning in cardiac myocytes. *Circ Res* 104: 879–886, 2009.
112. Rankin EB, Biju MP, Liu Q, Unger TL, Rha J, Johnson RS, Simon MC, Keith B, Haase VH. Hypoxia-inducible factor-2 (HIF-2) regulates hepatic erythropoietin in vivo. *J Clin Invest* 117: 1068–1077, 2007.
113. Rankin EB, Higgins DF, Walisser JA, Johnson RS, Bradfield CA, Haase VH. Inactivation of the arylhydrocarbon receptor nuclear translocator (Arnt) suppresses von Hippel-Lindau disease-associated vascular tumors in mice. *Mol Cell Biol* 25: 3163–3172, 2005.
114. Rondon LJ, MacMillan LA, Beckman BS, Goldberg MA, Schneider T, Bunn HF, Malter JS. Hypoxia up-regulates the activity of a novel erythropoietin mRNA binding protein. *J Biol Chem* 266: 16594–16598, 1991.
115. Rosenberger C, Mandriota S, Jurgensen JS, Wiesener MS, Horstrup JH, Frei U, Ratcliffe PJ, Maxwell PH, Bachmann S, Eckardt KU. Expression of hypoxia-inducible factor-1 α and -2 α in hypoxic and ischemic rat kidneys. *J Am Soc Nephrol* 13: 1721–1732, 2002.
116. Rosenberger C, Rosen S, Paliege A, Heyman SN. Pimonidazole adduct immunohistochemistry in the rat kidney: detection of tissue hypoxia. *Methods Mol Biol* 466: 161–174, 2009.
117. Rössler J, Stolze I, Frede S, Freitag P, Schweigerer L, Havers W, Fandrey J. Hypoxia-induced erythropoietin expression in human neuroblastoma requires a methylation free HIF-1 binding site. *J Cell Biochem* 93: 153–161, 2004.
118. Safran M, Kim WY, O'Connell F, Flippin L, Günzler V, Horner JW, Depinho RA, Kaelin WG Jr. Mouse model for noninvasive imaging of HIF prolyl hydroxylase activity: assessment of an oral agent that stimulates erythropoietin production. *Proc Natl Acad Sci USA* 103: 105–110, 2006.
119. Sakamoto T, Seiki M. Mint3 enhances the activity of hypoxia-inducible factor-1 (HIF-1) in macrophages by suppressing the activity of factor inhibiting HIF-1. *J Biol Chem* 284: 30350–30359, 2009.
120. Sanchez M, Galy B, Muckenthaler MU, Hentze MW. Iron-regulatory proteins limit hypoxia-inducible factor-2 α expression in iron deficiency. *Nat Struct Mol Biol* 14: 420–426, 2007.
121. Scandurro AB, Rondon LJ, Wilson RB, Tenenbaum SA, Garry RF, Beckman BS. Interaction of erythropoietin RNA binding protein with erythropoietin RNA requires an association with heat shock protein 70. *Kidney Int* 51: 579–584, 1997.
122. Schödel J, Klanke B, Weidemann A, Buchholz B, Bernhardt W, Bertog M, Amann K, Korbmacher C, Wiesener M, Warnecke C, Kurtz A, Eckardt KU, Willam C. HIF-prolyl hydroxylases in the rat kidney: physiologic expression patterns and regulation in acute kidney injury. *Am J Pathol* 174: 1663–1674, 2009.
123. Schofield CJ, Ratcliffe PJ. Oxygen sensing by HIF hydroxylases. *Nat Rev Mol Cell Biol* 5: 343–354, 2004.
124. Scortegagna M, Ding K, Zhang Q, Oktay Y, Bennett MJ, Bennett M, Shelton JM, Richardson JA, Moe O, Garcia JA. HIF-2 α regulates murine hematopoietic development in an erythropoietin-dependent manner. *Blood* 105: 3133–3140, 2005.
125. Selak MA, Armour SM, MacKenzie ED, Boulahbel H, Watson DG, Mansfield KD, Pan Y, Simon MC, Thompson CB, Gottlieb E. Succinate links TCA cycle dysfunction to oncogenesis by inhibiting HIF- α prolyl hydroxylase. *Cancer Cell* 7: 77–85, 2005.
126. Semenza GL, Koury ST, Nejfelt MK, Gearhart JD, Antonarakis SE. Cell-type-specific and hypoxia-inducible expression of the human erythropoietin gene in transgenic mice. *Proc Natl Acad Sci USA* 88: 8725–8729, 1991.
127. Seth P, Krop I, Porter D, Polyak K. Novel estrogen and tamoxifen induced genes identified by SAGE (Serial Analysis of Gene Expression). *Oncogene* 21: 836–843, 2002.
128. Shao Z, Zhang Y, Powell-Coffman JA. Two distinct roles for EGL-9 in the regulation of HIF-1-mediated gene expression in *Caenorhabditis elegans*. *Genetics* 183: 821–829, 2009.
129. Song YR, You SJ, Lee YM, Chin HJ, Chae DW, Oh YK, Joo KW, Han JS, Na KY. Activation of hypoxia-inducible factor attenuates renal injury in rat remnant kidney. *Nephrol Dial Transplant* 25: 77–85, 2010.
130. Steinhoff A, Pientka FK, Möckel S, Kettelhake A, Hartmann E, Köhler M, Depping R. Cellular oxygen sensing: Importins and exportins are mediators of intracellular localisation of prolyl-4-hydroxylases PHD1 and PHD2. *Biochem Biophys Res Commun* 387: 705–711, 2009.
131. Stiehl DP, Wirthner R, Köditz J, Spielmann P, Camenisch G, Wenger RH. Increased prolyl 4-hydroxylase domain proteins compensate for decreased oxygen levels. Evidence for an autoregulatory oxygen-sensing system. *J Biol Chem* 281: 23482–23491, 2006.
132. Takeda K, Aguila HL, Parikh NS, Li X, Lamothe K, Duan LJ, Takeda H, Lee FS, Fong GH. Regulation of adult erythropoiesis by prolyl hydroxylase domain proteins. *Blood* 111: 3229–3235, 2008.

133. **Takeda K, Cowan A, Fong GH.** Essential role for prolyl hydroxylase domain protein 2 in oxygen homeostasis of the adult vascular system. *Circulation* 116: 774–781, 2007.
134. **Takeda K, Ho VC, Takeda H, Duan LJ, Nagy A, Fong GH.** Placental but not heart defects are associated with elevated hypoxia-inducible factor α levels in mice lacking prolyl hydroxylase domain protein 2. *Mol Cell Biol* 26: 8336–8346, 2006.
135. **Tian H, Hammer RE, Matsumoto AM, Russell DW, McKnight SL.** The hypoxia-responsive transcription factor EPAS1 is essential for catecholamine homeostasis and protection against heart failure during embryonic development. *Genes Dev* 12: 3320–3324, 1998.
136. **Tian YM, Mole DR, Ratcliffe PJ, Gleadle JM.** Characterization of different isoforms of the HIF prolyl hydroxylase PHD1 generated by alternative initiation. *Biochem J* 397: 179–186, 2006.
137. **To KK, Huang LE.** Suppression of hypoxia-inducible factor 1 α (HIF-1 α) transcriptional activity by the HIF prolyl hydroxylase EGLN1. *J Biol Chem* 280: 38102–38107, 2005.
138. **van Wijk R, Sutherland S, Van Wesel AC, Huizinga EG, Percy MJ, Bierings M, Lee FS.** Erythrocytosis associated with a novel missense mutation in the *HIF2A* gene. *Haematologica*. In press.
139. **Wang GL, Jiang BH, Rue EA, Semenza GL.** Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O₂ tension. *Proc Natl Acad Sci USA* 92: 5510–5514, 1995.
140. **Warnecke C, Zaborowska Z, Kurreck J, Erdmann VA, Frei U, Wiesener M, Eckardt KU.** Differentiating the functional role of hypoxia-inducible factor (HIF)-1 α and HIF-2 α (EPAS-1) by the use of RNA interference: erythropoietin is a HIF-2 α target gene in Hep3B and Kelly cells. *FASEB J* 18: 1462–1464, 2004.
141. **Wenger RH.** Cellular adaptation to hypoxia: O₂-sensing protein hydroxylases, hypoxia-inducible transcription factors, and O₂-regulated gene expression. *FASEB J* 16: 1151–1162, 2002.
142. **Wenger RH, Camenisch G, Stiehl DP, Katschinski DM.** HIF prolyl-4-hydroxylase interacting proteins: consequences for drug targeting. *Curr Pharm Des* 15: 3886–3894, 2009.
143. **Wenger RH, Kvietikova I, Rolfs A, Camenisch G, Gassmann M.** Oxygen-regulated erythropoietin gene expression is dependent on a CpG methylation-free hypoxia-inducible factor-1 DNA-binding site. *Eur J Biochem* 253: 771–777, 1998.
144. **Wenger RH, Stiehl DP, Camenisch G.** Integration of oxygen signaling at the consensus HRE. *Sci STKE* 2005: re12, 2005.
145. **Wiesener MS, Seyfarth M, Warnecke C, Jurgensen JS, Rosenberger C, Morgan NV, Maher ER, Frei U, Eckardt KU.** Paraneoplastic erythrocytosis associated with an inactivating point mutation of the von Hippel-Lindau gene in a renal cell carcinoma. *Blood* 99: 3562–3565, 2002.
146. **Wirthner R, Balamurugan K, Stiehl DP, Barth S, Spielmann P, Oehme F, Flamme I, Katschinski DM, Wenger RH, Camenisch G.** Determination and modulation of prolyl-4-hydroxylase domain oxygen sensor activity. *Methods Enzymol* 435: 43–60, 2007.
147. **Wong W, Goehring AS, Kapiloff MS, Langeberg LK, Scott JD.** mAKAP compartmentalizes oxygen-dependent control of HIF-1 α . *Sci Signal* 1: ra18, 2008.
148. **Zhao S, Lin Y, Xu W, Jiang W, Zha Z, Wang P, Yu W, Li Z, Gong L, Peng Y, Ding J, Lei Q, Guan KL, Xiong Y.** Glioma-derived mutations in IDH1 dominantly inhibit IDH1 catalytic activity and induce HIF-1 α . *Science* 324: 261–265, 2009.

